Center for Crop Diversification Crop Profile CCD-CP-62

High Tunnel Tomatoes

Cheryl Kaiser¹ and Matt Ernst²

Introduction

High tunnels, also known as hoop houses, are simple polyethylene-covered unheated structures. Tunnels can be covered with one or two sheets of plastic; those covered with two have an air layer in between, thus offering better insulation and, consequently, more cold protection (and wind protection). High tunnels are used to extend the growing season earlier into spring and later into fall. Determinate and indeterminate tomatoes (Lycopersicon esculentum) can be successfully grown in this production system, yielding a potentially profitable "out of season" crop. Other advantages to high tunnel production include: higher marketable yields, improved fruit quality, fewer culls, and a reduction in pesticide applications.

University of Kentucky College of Agriculture, Food and Environment

Cooperative Extension Service

Marketing and Market Outlook

Growers able to provide the earliest locally grown tomatoes can often demand a premium price as customers are eager for that first truly vine-ripened tomato. High tunnel tomato production can yield a crop four to five weeks ahead of field-grown tomatoes, and producers who capture the early market may gain loyal season-long customers. Similarly, the scarcity of locally grown late-season tomatoes can provide a profitable market for growers able to extend production well into fall. Extending the season has the potential of spreading out cash flow, increasing overall farm profits, and gaining new customers.

Marketing avenues include roadside stands, farmers

markets, restaurants, locally owned community grocers. supported agriculture (CSA), produce auctions, institutions (such as schools and hospitals), and wholesale grocery and DIVERSIFICATION foodservice markets.





Production considerations

Growing environment

High tunnels are passively solar-heated. Nighttime low temperatures within a high tunnel can be a few degrees warmer than outside temperatures. Row covers used in conjunction with the high tunnel will provide additional cold and frost protection. As an alternative, a heater can be used when unexpected drops in temperature occur. However, due to rising energy costs, the use of heaters should generally be limited to short durations. A grower who has received NRCS EQIP funds to purchase a high tunnel should check his/ her state's NRCS High Tunnel System Conservation Practice Standard to make sure heaters are allowed. NRCS allows the permanent installation of heaters in Kentucky. Such additions should be included in the manufacturer's design and recommendations.

> During sunny days, temperatures in the tunnels can reach 30° F to 40° F higher than outside temperatures, necessitating that tunnels be adequately vented. Lacking the automated system of more

¹Cheryl Kaiser is a former Extension Associate with the Center for Crop Diversification. ²Matt Ernst is an independent contractor with the Center for Crop Diversification.

Cooperative Extension Service | Agriculture and Natural Resources | Family and Consumer Sciences | 4-H Youth Development | Community and Economic Development

elaborate greenhouses, high tunnels are ventilated by manually rolling up the sidewalls in the morning and closing them at night. Orienting the houses perpendicular to the prevailing wind on the farm will help facilitate air movement. It is also important that the sidewalls be high enough to allow for adequate cross ventilation. High tunnels should not be closed at night if the ambient temperature is greater than 55° F. A shade cloth may be needed in early summer to prevent sunburn of tomato fruit.

Because tunnels prevent natural rainfall from reaching plants, drip irrigation is essential for providing the large volume of water this crop requires for quality fruit production. The irrigation system can also be

used to provide fertilization during the season in the high tunnel.

Plant selection

Growers interested in an early tomato should crop select a determinate cultivar. Determinates produce a heavy flush of tomatoes over a period of about four to six weeks. With proper planning, their production will end about the time field tomatoes begin to bear. On the other hand, indeterminate tomatoes produce a more sustained harvest over a longer period of time. Some producers will grow indeterminate cultivars early in the spring and keep them productive throughout the

summer and into late fall when prices begin to rise again.

Heirloom tomatoes are also a good crop for this production system because they are less prone to cracking and have fewer disease problems in the tunnel than in the field.

Site selection

Select a level, well-drained fertile location for tomato production. Site selection is more critical when the high tunnel is built to be a permanent structure. Once permanent tunnels are in place, amending the soil each year can present a challenge. A small tractor with a tiller can be driven into tunnels with hinged endwalls. However, if the endwalls are fixed, soil preparation will need to be accomplished by hand or with a small power tiller. The soil in moveable tunnels can be worked prior to erecting the structure.

Moveable high tunnels can be relocated to a new site each season to facilitate crop rotation and to avoid nutrient depletion of the soil. Rotating moveable high tunnels to different locations also helps to prevent the buildup of soil-borne diseases and/or fertilizer salts. Often tunnels that remain in the same location, producing the same crop for a number of years, will have high levels of soil pathogens, leading to progressively high disease loads and crop losses.

> Fertilizer salt build-up can also be avoided when the tunnel is moveable. If the tunnel is not moved, it is important to remove the plastic every third or fourth year to allow salts to leach from the soil.

> Raised beds of at least 6 inches will promote earliness by enhancing soil warming while also improving soil drainage. On the other hand, growers who produce greens in the same site during the winter months may decide to forgo raised beds; soils in raised beds do not stay as warm for winter crop production as soil-level beds. Using black plastic mulch will increase soil temperature, provide weed

control, and reduce water loss. If organic mulch is preferred to plastic, applications should be made after the soil has warmed because some organic mulches have a tendency to actually lower soil temperatures.

Planting and crop support

Healthy 4- to 5-week-old seedlings can be transplanted to the tunnel once the soil has warmed to 60° F at a 2-inch depth. For permanent high tunnels the soil stays relatively warm, which means that growers in Zone 6 are often able to plant four to five weeks earlier than normal in the spring. Fall plantings can be made in early August for mid-November to mid-December harvests.



Many tomato cultivars, including determinates, grow considerably taller in the tunnel environment than they would in the field. Crop support is very important in a high tunnel for many of the same reasons it is used in the field: better light penetration, improved air circulation, more plants per square foot, and fewer fruit in contact with the ground. Options include cages, staking, stake and weave, and trellising; some methods are more labor intensive and/or costly than others. Trellising to the tunnel roof is not advisable unless the frame has been built to support the heavy weight of fruit-laden plants. However, a separate support system can be constructed inside the high tunnel if trellising is preferred. A 20-foot by 96-foot tunnel can accommodate from 125 to 300 tomato plants depending on spacing, bed size, width of walkways, and height of sidewalls.

Pruning and pollinating

Suckers (axillary shoots) should be pruned to hasten earliness and improve air circulation. Prune only those below the first flower cluster that are less than 4 inches long. Pollination of tunnel-produced tomatoes is not adversely affected as long as there is good air movement through the structure. If the tunnel sidewalls need to be kept down for extended periods of time, it may be necessary to promote pollination by vibrating plants, using a leaf blower, or by tapping the crop support system (i.e. stakes or cages).

Pest management

Due to the unique environment within tunnels, growers may encounter different disease problems in the tunnel than in the field. Because the tunnel excludes rainfall, the foliage tends to stay dry, resulting in fewer disease problems from pathogens that are spread by rain splash or require leaf-wetness for infection. However, due to the limited movement of air in a tunnel, those diseases favored by high humidity can increase in severity. Because of potential virus problems, bedding plants and other potted ornamentals should not be planted in the same greenhouse as tomatoes.

Due to the irrigation needs of tunnel-grown tomatoes, blossom end rot, a calcium deficiency caused by irregular watering, can be a problem for some growers. Insect pests in tunnels differ as well. While the high tunnel presents a barrier to some insect pests (e.g. the sphinx moth), it is an ideal environment for others, particularly whiteflies and aphids. Typically, any insect pest that is a problem for greenhouses will be a problem in high tunnels. An insect exclusion screen can be placed over the sidewall and gable vents to prevent insect entry into the high tunnel.

Frequent scouting to monitor insect populations and disease incidence is essential. Due to the relatively high density of plants in tunnels, pests and diseases can spread very quickly.

Harvest and storage

Tomatoes for fresh market are harvested when fully vine-ripe. Harvesting early in the morning before the tunnel heats up is best for both the picker and the tomato. To avoid crushing, fruit should not be packed more than two layers thick. Determinate plants can yield from 10 to 25 pounds of tomatoes per plant over a four-week period. High tunnel tomatoes generally produce fewer culls than field-grown plants.

Labor requirements

Labor figures provided here are for high tunnel tomato production in a single 96-foot x 20-foot house. Site preparation, which includes soil preparation and getting the tunnel ready, is estimated at 10 to 12 hours. Approximately three hours will be needed for planting and two hours for staking. Labor during production (15 to 30 hours) mainly involves such labor-intensive activities as pruning, hand spraying or weeding. Harvesting and packing is expected to demand about 40 hours. In addition, high tunnels will require daily labor to manually raise and lower sidewalls for temperature and ventilation management (10 hours). These structures could also require monitoring during heavy storms. Additional labor may be necessary for producers marketing their own tomatoes.

Economic considerations

Initial investments include high tunnel construction, land preparation, purchase of seed or transplants, and installation of an irrigation system. Additional start-up costs can include black plastic mulch, row covers, an inflation fan, and a crop support system.

High tunnels are a relatively inexpensive way to extend the growing season. Excluding labor, the approximate materials cost of high tunnel construction is \$1.30 to \$1.50 per square foot. Site preparation and high tunnel systems, like ventilation fans and heating, can add about the same amount per square foot. High

tunnel operators in Kentucky reported high tunnel construction costs during winter 2017 at \$2.50 to \$3 per square foot, plus the value of some operator labor. Compared to the cost of a traditional productionready greenhouse (\$8 to \$30 per square foot), the high tunnel requires little capital investment. High tunnel production does represent a much greater investment than growing produce in the field.

The following 2016 budget information is based on production of 200 plants in a 96-foot by 20-foot high tunnel. Production costs for irrigated tomatoes are estimated at \$760. Harvest and marketing costs, including 20 hours of hired labor, are estimated at \$630. Total expenses, including both variable and a \$550 annual fixed cost, would come to approximately \$2,050. Assuming tomato gross returns of \$3,300 and an average wage rate of \$10/hour, returns to land, capital and management would be \$850. These estimates assume a price of \$2 per pound for 75 percent of a 2,000-pound yield. Total returns per high tunnel can be increased by raising tomato planting populations or utilizing unused space within the high tunnel for other crop production. Higher returns may also be obtained by marketing specialty varieties as well as selling early season tomatoes at higher prices.

Selected Resources

• How to Build a High Tunnel (University of Kentucky) <u>http://www.uky.edu/hort/sites/www.uky.edu.hort/files/documents/hightunnel.pdf</u>

• An IPM Scouting Guide for Common Problems of High Tunnel and Greenhouse Vegetable Crops in Kentucky (University of Kentucky, 2016) http://www2.ca.uky.edu/agcomm/pubs/ID/ID235/ ID235.pdf • Economics of High Tunnel Vegetable and Strawberry Production in the Central Midwest (West Virginia University, 2007) 6.7 MB file <u>https://www2.ag.purdue.edu/hla/fruitveg/</u> <u>Presentations/econ_summer_crop.pdf</u>

• High Tunnel Tomato Production Basics (Penn State University, 2017)

https://extension.psu.edu/high-tunnel-tomatoproduction-basics?

Tomatoes for the High Tunnel (Cornell University, 2014) <u>https://rvpadmin.cce.cornell.edu/uploads/</u>
<u>doc_201.pdf</u>

• High Tunnel Tomato Production (Utah State University, 2010) <u>http://extension.usu.edu/</u> files/publications/publication/Horticulture_ <u>HIghTunnels_2010-03pr.pdf</u>

• Minnesota High Tunnel Production Manual (University of Minnesota Extension, 2012) <u>http://www.extension.umn.edu/garden/fruit-</u> <u>vegetable/docs/high-tunnel-manual-2012.pdf</u>

• Warm Season Vegetables and Melons (Hightunnels.org - Kansas State Research and Extension, University of Missouri Extension, and University of Nebraska Cooperative Extension) http://hightunnels.org/category/for-growers/growingin-high-tunnels/warm-season-vegetables-melons/

• High Tunnel Production (Virginia Tech) http://www.hort.vt.edu/ghvegetables/hightunnel.html

Suggested Citation:

Kaiser, C. and M. Ernst. (2017). *High Tunnel Tomatoes*. CCD-CP-62. Lexington, KY: Center for Crop Diversification, University of Kentucky College of Agriculture, Food and Environment. Available: <u>http://www.uky.edu/ccd/sites/www.</u> uky.edu.ccd/files/hightunneltomatoes.pdf

Reviewed by Lewis Jett, Extension Specialist, West Virginia University Photos courtesy of Alexandra Williams, Technical Expert, Syngenta (Page 1), and Krista Jacobsen, Assistant Professor, University of Kentucky (Page 2)

September 2017

For additional information, contact your local County Extension agent

Educational programs of Kentucky Cooperative Extension serve all people regardless of economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender expression, pregnancy, marital status, genetic information, age, veteran status, or physical or mental disability.